


CT/EP 91/00620

Sheet number 4

<b>Box No. VI PRIORITY CLAIM (IF ANY).</b> The priority of the following earlier application(s) is hereby claimed:			
Country (country in which it was filed if national application; one of the countries for which it was filed if regional or international application)	Filing Date (day, month, year) <div style="text-align: center; font-size: 1.2em;">(02.04.90)</div>	Application No. <div style="text-align: center;">90810262.7</div>	Office of Filing (fill in only if the earlier application is an international application or a regional application) <div style="text-align: center; font-size: 1.2em;">EP</div>
(1) [Europe] DE	April 2, 1990		
(2)			
(3)			
(Letter codes may be used to indicate country and/or Office of filing)			
When the earlier application was filed with the Office which, for the purposes of the present international application, is the receiving Office, the applicant may, <i>against payment of the required fee</i> , ask the following: <input type="checkbox"/> the receiving Office is hereby requested to prepare and transmit to the International Bureau a certified copy of the above-mentioned earlier application/of the earlier applications identified above by the numbers (insert the applicable numbers)			
<b>Box No. VII EARLIER SEARCH (IF ANY).</b> Fill in where a search (international, international-type or other) by the International Searching Authority has already been requested (or completed) and the said Authority is now requested to base the international search, to the extent possible, on the results of the said earlier search. Identify such search or request either by reference to the relevant application (or the translation thereof) or by reference to the search request.			
International application number or number and country (or regional Office) of other application: EP 90810262.7		International/regional/national filing date: April 2, 1990	
Date of request for search:		Number (if available) given to search request:	
<b>Box No. VIII SIGNATURE OF APPLICANT(S) OR AGENT</b>			
 Blaise Doussé & Jean-Paul Savoye CONSEILS EN PROPRIÉTÉ INDUSTRIELLE 7, rue de Diderot CH-1227 Carouge (Genève-Suisse)			
If the present Request form is signed on behalf of any applicant by an agent, a separate power of attorney appointing the agent and signed by the applicant is required. If in such case it is desired to make use of a general power of attorney (deposited with the receiving Office), a copy thereof must be attached to this form.			
<b>Box No. IX CHECK LIST (To be filled in by the Applicant)</b>		This international application as filed is accompanied by the items checked below:	
This international application contains the following number of sheets:		1. <input type="checkbox"/> separate signed power of attorney 2. <input type="checkbox"/> copy of general power of attorney 3. <input type="checkbox"/> priority document(s) (see Box No. VI) 4. <input type="checkbox"/> receipt of the fees paid or revenue stamps 5. <input type="checkbox"/> cheque for the payment of fees 6. <input type="checkbox"/> request to charge deposit account 7. <input checked="" type="checkbox"/> other document (specify) copie of the European Search Report	
1. request _____ 4 sheets 2. description _____ 21 sheets 3. claims _____ 4 sheets 4. abstract _____ 1 sheet 5. drawings _____ sheets <div style="text-align: right;">Total 30 sheets</div>	Figure number _____ of the drawings (if any) is suggested to accompany the abstract for publication.		
(The following is to be filled in by the receiving Office)			
1. Date of actual receipt of the purported international application:		02 APR 1991 (02.04.91)	
2. Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application:			
3. Date of timely receipt of the required corrections under Article 11 of the PCT:			
4. Drawings <input type="checkbox"/> Received <input checked="" type="checkbox"/> No Drawings			
(The following is to be filled in by the International Bureau)			
Date of receipt of the record copy:		25 APRIL 1991 (25.04.91)	

Europäisch  
Patentamt

European Patent  
Office

Office européen  
des brevets

PCT/EP 91/00620

27 Rec'd PCT/PTO 04 NOV 91

PRIORITY DOCUMENT



REC'D 29 APR 1991  
WIPO PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Den Haag, den  
The Hague,  
La Haye, le

02.04.91

Der Präsident des Europäischen Patentamts  
Im Auftrag  
For the President of the European Patent Office  
Le Président de l'Office européen des brevets  
p. o.

P. J. MASSAAR

Patentanmeldung Nr. 90810262.7  
Patent application no.  
Demande de brevet n°

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**



Anmeldung Nr.: 90810262.7  
Application no.:  
Demande n°:

Anmeldetag:  
Date of filing: 02.04.1990  
Date de dépôt:

Anmelder: SINTETICA SA  
Applicant(s): Mendrisio - Switzerland  
Demandeur(s):

Bezeichnung der Erfindung: *Stable microbubbles suspensions injectable into living organisms.*  
Title of the invention:  
Titre de l'invention:

In Anspruch genommene Priorität(en)  
Priority(ies) claimed  
Priorité(s) revendiquée(s)

Staat: ---  
State:  
Pays:

Tag:  
Date: ---  
Date:

Aktenzeichen:  
File no. ---  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent Classification: A 61 K 49/00, A 61 B 8/00  
Classification internationale des brevets:

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at the date of filing:  
Etats contractants désignés lors du dépôt:

AT-BE-CH-LI-DE-ES-FR-GB-GR-IT-LU-NL-SE/DK

Bemerkungen:  
Remarks:  
Remarques:

## STABLE MICROBUBBLES SUSPENSIONS INJECTABLE INTO LIVING ORGANISMS

The present invention concerns media adapted for injection into living bodies, e.g. for the purpose of ultrasonic echography and, more particularly, injectable liquid compositions comprising microbubbles of air or physiologically acceptable gases as stable dispersions or suspensions in an aqueous liquid carrier. These compositions are mostly usable as contrast agents in ultrasonic echography to image the inside of blood-stream vessels and other cavities of living beings, e.g. human patients and animals. Other uses however are also contemplated as disclosed hereafter.

The invention also comprises dry compositions which, upon admixing with an aqueous carrier liquid, will generate the foregoing sterile suspension of microbubbles thereafter usable as contrast agent for ultrasonic echography and other purposes.

It is well known that gas microbubbles suspended in a liquid are exceptionally efficient ultrasound reflectors. Hence injecting into the blood-stream of living bodies suspensions of gas microbubbles (in the range of 0.5 to 10  $\mu\text{m}$ ) in a carrier liquid will strongly reinforce ultrasonic echography imaging, thus aiding in the visualization of internal organs. Imaging of vessels and internal organs can strongly help in medical diagnosis, for instance for the detection of cardiovascular and other diseases.

The generation of suspensions of gas or air filled microbubbles (0.5 - 50  $\mu\text{m}$ ) in an injectable carrier liquid can follow various routes. For instance in DE-A- 3529195 (Max-Planck Gesell.), there is disclosed a technique in which an aqueous emulsified mixture containing a water soluble polymer, an oil and mineral salts is forced back and forth, together with a small amount of air, from one syringe into another through a small opening. Here, mechanical forces are responsible for the formation of bubbles in the liquid.

M.W. Keller et al. (J. Ultrasound Med. 5 (1986), 439-8)

have reported subjecting to ultrasonic cavitation under atmospheric pressure solutions containing high concentrations of solutes such as dextrose, Renografin-76, Iopamidol (an X-ray contrast agent), and the like. There the air is driven into the solution by the energy of cavitation.

Other techniques rely on the shaking of a carrier liquid in which air containing microparticles have been incorporated, said carrier liquid usually containing, as stabilizers, viscosity enhancing agents, e.g. water soluble polypeptides or carbohydrates and/or surfactants. It is effectively admitted that the stability of the microbubbles against decay or escape to the atmosphere is controlled by the viscosity and surface properties of the carrier liquid. The air or gas in the microparticles can consist of inter-particle or intra-crystalline entrapped gas, as well as surface adsorbed gas, or gas produced by reactions with the carrier liquid, usually aqueous. All this is fully described for instance in EP-A- 52.575 (Ultra Med. Inc.) in which there are used aggregates of 1 - 50  $\mu\text{m}$  particles of carbohydrates (e.g. galactose, maltose, sorbitol, gluconic acid, sucrose, glucose and the like) in aqueous solutions of glycols or polyglycols, or other water soluble polymers.

Also, in EP-A- 123.235 and 122.624 (Schering) use is made of air trapped in solids. For instance, 122.624 claims a liquid carrier contrast composition for ultrasonic echography containing microparticles of a solid surfactant, the latter being optionally combined with microparticles of a non-surfactant. As explained in this document, the formation of air bubbles in the solution results from the release of the air adsorbed on the surface of the particles, or trapped within the particle lattice, or caught between individual particles, this being so when the particles are agitated with the liquid carrier.

EP-A- 131.540 (Schering) also discloses a method in which a stabilized carrier liquid, e.g. a physiological aqueous solution of salt, or a solution of a sugar like maltose, dextrose, lactose or galactose, without viscosity enhancer, is mixed with microparticles (in the 0.1 to 1  $\mu\text{m}$  range) of the same sugars containing entrapped air.

In order that the suspension of bubbles can develop within

the liquid carrier, the foregoing documents recommend that both liquid and solid components be violently agitated together under sterile conditions; the agitation of both components together is performed for a few seconds and the suspension must then be used immediately, i.e. it should be injected within 5 - 10 minutes for echographic measurements; this indicates that the bubbles in the suspensions are not longlived.

In US-A- 4,466,442 (Schering), there is disclosed a series of different techniques for producing suspensions of gas microbubbles in a liquid carrier using (a) a solution of a tenside (surfactant) in a carrier liquid (aqueous) and (b) a solution of a viscosity enhancer as stabilizer. For generating the bubbles, the techniques used there include forcing at high velocity a mixture of (a), (b) and air through a small aperture; or injecting (a) into (b) shortly before use together with a physiologically acceptable gas; or adding an acid to (a) and a carbonate to (b), both components being mixed together just before use and the acid reacting with the carbonate to generate CO<sub>2</sub> bubbles; or adding an over-pressurized gas to a mixture of (a) and (b) under storage, said gas being released into microbubbles at the time when the mixture is used for injection.

The tensides used in component (a) of US-A- 4,466,442 comprise lecithins; esters and ethers of fatty acids and fatty alcohols with polyoxyethylene and polyoxyethylated polyols like sorbitol, glycols and glycerol, cholesterol; and polyoxy-ethylene-polyoxypropylene polymers. The viscosity raising and stabilizing compounds include for instance mono- and polysaccharides (glucose, lactose, sucrose, dextran, sorbitol); polyols, e.g. glycerol, polyglycols; and polypeptides like proteins, gelatin, oxypolygelatin, plasma protein and the like.

In a typical preferred example of this document, equivalent volumes of (a) a 0.5% by weight aqueous solution of Pluronic® F-68 (a polyoxypropylene-polyoxyethylene polymer) and (b) a 10% lactose solution are vigorously shaken together under sterile conditions (closed vials) to provide a suspension of microbubbles ready for use as an ultrasonic contrast agent and lasting for at least 2 minutes. About 50% of the bubbles had a size below 50 µm.

Although the achievements of the prior art have merit, they suffer from several drawbacks which strongly limit their practical use by doctors and hospitals, namely their relatively short life-span (which makes test reproducibility difficult), relative low initial bubble concentration (the number of bubbles rarely exceeds  $10^4 - 10^5$  bubbles/ml and the count decreases rapidly with time) and poor reproducibility of the initial bubble count from test to test (which also makes comparisons difficult). Also it is admitted that for efficiently imaging certain organs, e.g. the left heart, bubbles smaller than  $50 \mu\text{m}$ , preferably in the range of  $0.5-10 \mu\text{m}$ , are required.

Furthermore, the compulsory presence of solid microparticles or high concentrations of electrolytes and other relatively inert solutes in the carrier liquid may be undesirable physiologically in some cases. Finally, the suspensions are totally unstable under storage and cannot be marketed as such; hence great skill is required to prepare the microbubbles at the right moment just before use.

Of course there exists stable suspensions of microcapsules, i.e. microballoons with a solid, air-sealed rigid polymeric membrane which perfectly resist for long storage periods in suspension, which have been developed to remedy this shortcoming (see for instance K.J. Widder, EP-A- 324.938); however the properties of microcapsules in which a gas is entrapped inside solid membrane vesicles essentially differ from that of the gas microbubbles of the present invention and belong to a different kind of art; for instance while the gas microbubbles discussed here will simply escape or dissolve in the blood-stream when the stabilizers in the carrier liquid are excreted or metabolized, the solid polymer material forming the walls of the aforementioned micro-balloons must eventually be disposed of by the organism being tested which may impose a serious afterburden upon it. Also capsules with solid, non-elastic membrane may break irreversibly under variations of pressure.

The composition of the present invention, as defined in claim 1, fully remedies the aforementioned pitfalls.

The term "lamellar form" defining the condition of at least a portion of the surfactant or surfactants of the present compo-

sition indicates that the surfactants, in strong contrast with the microparticles of the prior art (for instance EP-A-123.235), are in the form of thin films involving one or more molecular layers (in laminate form). Converting film forming surfactants into lamellar form can easily be done for instance by high pressure homogenization or by sonication under acoustical or ultrasonic frequencies. In this connection, it should be pointed out that the existence of liposomes is a well known and useful illustration of cases in which surfactants, more particularly lipids, are in lamellar form.

Liposome solutions are aqueous suspensions of microscopic vesicles, generally spherically shaped, which hold substances encapsulated therein. These vesicles are usually formed of one or more concentrically arranged molecular layers (lamellae) of amphipatic compounds, i.e. compounds having a lipophobic hydrophilic moiety and a lipophilic hydrophobic moiety. See for instance "Liposome Methodology", Ed. L.D. Leserman et al, Inserm 136, 2-8 May 1982). Many surfactants or tensides, including lipids, particularly phospholipids, can be laminarized to correspond to this kind of structure. In this invention, one preferably uses the lipids commonly used for making liposomes, for instance the lecithins and other tensides disclosed in more detail hereafter, but this does in no way preclude the use of other surfactants provided they can be formed into layers or films.

Practically, to achieve the suspensions of microbubbles according to the invention, one may start with liposomes suspensions or solutions prepared by any technique reported in the prior art, with the obvious difference that in the present case the liposomic vesicles are preferably "unloaded", i.e. they do not need to keep encapsulated therein any foreign material as is normally the object of classic liposomes. Hence, preferably, the liposomes of the present invention will contain an aqueous phase identical or similar to the aqueous phase of the solution itself. Then air or a gas is introduced into the liposome solution so that a suspension of microbubbles will form, said suspension being stabilized by the presence of the surfactants in lamellar form. Notwithstanding, the material making the liposome walls



can be modified within the scope of the present invention, for instance by covalently grafting thereon foreign molecules designed for specific purposes as will be explained later.

The preparation of liposome solutions has been abundantly discussed in many publications, e.g. US-A- 4,224,179 and WO-A-88/09165 and all citations mentioned therein. This prior art is used here as reference for exemplifying the various methods suitable for converting film forming tensides into lamellar form. Another basic reference by M.C. Woodle and D. Papahadjopoulos is found in "Methods in Enzymology" 171 (1989), 193.

For instance, in a method disclosed in D.A. Tyrrell et al, Biochimica & Biophysica Acta 457 (1976), 259-302, a mixture of a lipid and an aqueous liquid carrier is subjected to violent agitation and thereafter sonicated at acoustic or ultrasonic frequencies at room or elevated temperature. In the present invention, it has been found that sonication without agitation is convenient. Also, an apparatus for making liposomes, a high pressure homogenizer such as the Microfluidizer<sup>®</sup>, which can be purchased from Microfluidics Corp., Newton, MA 02164 USA, can be used advantageously. Large volumes of liposome solutions can be prepared with this apparatus under pressures which can reach 600-1200 bar.

According to the teaching of GB-A- 2,134,869 (Squibb), microparticles (10  $\mu$ m or less) of a hydrosoluble carrier solid (NaCl, sucrose, lactose and other carbohydrates) are coated with an amphipatic agent; the dissolution of the coated carrier in an aqueous phase will yield liposomic vesicles. In GB-A- 2,135,647 insoluble particles, e.g. glass or resin microbeads are coated by moistening in a solution of a lipid in an organic solvent followed by removal of the solvent by evaporation. The lipid coated microbeads are thereafter contacted with an aqueous carrier phase, whereby liposomic vesicles will form in that carrier phase.

The introduction of air or gas into a liposome solution in order to form therein a suspension of microbubbles can be effected by usual means, inter alia by injection, that is, forcing said air or gas through tiny orifices into the liposome

solution, or simply dissolving the gas in the solution by applying pressure and thereafter suddenly releasing the pressure. Another way is to agitate or sonicate the liposome solution in the presence of air or an entrappable gas. Also one can generate the formation of a gas within the solution of liposomes itself, for instance by a gas releasing chemical reaction, e.g. decomposing a dissolved carbonate or bicarbonate by acid. The same effect can be obtained by dissolving under pressure a low boiling liquid, for instance butane, in the aqueous phase and thereafter allowing said liquid to boil by suddenly releasing the pressure.

Notwithstanding, an advantageous method is to contact the dry surfactant in lamellar or thin film form with air or an adsorbable or entrappable gas before introducing said surfactant into the liquid carrier phase. In this regard, the method can be derived from the technique disclosed in GB-A-2,135,647, i.e. solid microparticles or beads are dipped in a solution of a film forming surfactant (or mixture of surfactants) in a volatile solvent, after which the solvent is evaporated and the beads are left in contact with air (or an adsorbable gas) for a time sufficient for that air to become superficially bound to the surfactant layer. Thereafter, the beads coated with air filled surfactant are put into a carrier liquid, usually water with or without additives, whereby air bubbles will develop within the liquid by gentle mixing, violent agitation being entirely unnecessary. Then the solid beads can be separated, for instance by filtration, from the microbubble suspension which is remarkably stable with time.

Needless to say that, instead of insoluble beads or spheres, one may use as supporting particles water soluble materials like that disclosed in GB-A- 2,134,869 (carbohydrates or hydrophilic polymers), whereby said supporting particles will eventually dissolve and final separation of a solid becomes unnecessary. Furthermore in this case, the material of the particles can be selected to eventually act as stabilizer or viscosity enhancer wherever desired.

In a variant of the method, one may also start with dehydrated liposomes, i.e. liposomes which have been prepared

normally by means of conventional techniques in the form of aqueous solutions and thereafter dehydrated by usual means, e.g. such as disclosed in US-A- 4,229,360 also incorporated herein as reference. One of the methods for dehydrating liposomes recommended in this reference is freeze-drying (lyophilization), i.e. the liposome solution is frozen and dried by evaporation (sublimation) under reduced pressure. Prior to effecting freeze-drying, a hydrophilic stabilizer compound is dissolved in the solution, for instance a carbohydrate like lactose or sucrose or a hydrophilic polymer like dextran, starch, PVP, PVA and the like. This is useful in the present invention since such hydrophilic compounds also aid in homogenizing the microbubbles size distribution and enhance stability under storage. Actually making very dilute aqueous solutions (0.1 - 10% by weight) of freeze-dried liposomes stabilized with, for instance, a 5:1 to 10:1 weight ratio of lactose to lipid enables to produce aqueous microbubbles suspensions counting  $10^8$  -  $10^9$  microbubbles/ml (size distribution mainly 0.5 - 10  $\mu$ m) which are stable for at least a month (and probably much longer) without significant observable change. And this is obtained by simple dissolution of the air-stored dried liposomes without shaking or any violent agitation. Furthermore, the freeze-drying technique under reduced pressure is very useful because it permits, after drying, to restore the pressure above the dried liposomes with any entrappable gas, i.e. nitrogen, CO<sub>2</sub>, argon, methane, freon, etc., whereby after dissolution of the liposomes processed under such conditions suspensions of microbubbles containing the above gases are obtained.

Microbubbles suspensions formed by applying gas pressure on a dilute solution of laminated lipids in water (0.1 - 10% by weight) and thereafter suddenly releasing the pressure have an even higher bubble concentration, e.g. in the order of  $10^{10}$  -  $10^{11}$  bubbles/ml. However, the average bubble size is somewhat above 10  $\mu$ m, e.g. in the 10 - 50  $\mu$ m range. In this case, bubble size distribution can be narrowed by centrifugation and layer decantation.

The tensides or surfactants which are convenient in this invention can be selected from all amphipatic compounds capable

of forming stable films in the presence of water and gases. The preferred surfactants include the lecithins (phosphatidyl-choline) and other phospholipids, inter alia phosphatidic acid (PA), phosphatidyl-inositol phosphatidyl-ethanolamine (PE), phosphatidyl-serine (PS), phosphatidyl-glycerol (PG), cardiolipin (CL), sphingomyelins, the plasmogens, the cerebrosides, etc. Examples of suitable lipids are the phospholipids in general, for example, natural lecithins, such as egg lecithin or soya bean lecithin, or synthetic lecithins such as saturated synthetic lecithins, for example, dimyristoyl phosphatidyl choline, dipalmitoyl phosphatidyl choline or distearoyl phosphatidyl choline or unsaturated synthetic lecithins, such as dioleoyl phosphatidyl choline or dilinoleoyl phosphatidyl choline, with egg lecithin or soya bean lecithin being preferred. Additives like cholesterol and other substances (see below) can be added to one or more of the foregoing lipids in proportions ranging from zero to 50% by weight.

Such additives may include other surfactants that can be used in admixture with the film forming surfactants and most of which are recited in the prior art discussed in the introduction of this specification. For instance, one may cite free fatty acids, esters of fatty acids with polyoxyalkylene compounds like polyoxypropylene glycol and polyoxyethylene glycol; ethers of fatty alcohols with polyoxyalkylene glycols; esters of fatty acids with polyoxyalkylated sorbitan; soaps; glycerol-polyalkylene stearate; glycerol-polyoxyethylene ricinoleate; homo- and copolymers of polyalkylene glycols; polyethoxylated soya-oil and castor oil as well as hydrogenated derivatives; ethers and esters of sucrose or other carbohydrates with fatty acids, fatty alcohols, these being optionally polyoxyalkylated; mono-, di- and triglycerides of saturated or unsaturated fatty acids; glycerides of soya-oil and sucrose. The amount of the non-film forming tensides or surfactants can be up to 50% by weight of the total amount of surfactants in the composition but is preferably between zero and 30%.

The total amount of surfactants relative to the aqueous carrier liquid is best in the range of 0.01 to 25% by weight but quantities in the range 0.5 - 5% are advantageous because one

always tries to keep the amount of active substances in an injectable solution as low as possible, this being to minimize the introduction of foreign materials into living beings even when they are harmless and physiologically compatible.

Further optional additives to the surfactants include:

a) substances which are known to provide a negative charge on liposomes, for example, phosphatidic acid, phosphatidyl-glycerol or dicetyl phosphate;

b) substances known to provide a positive charge, for example, stearyl amine, or stearyl amine acetate;

c) substances known to affect the physical properties of the lipid films in a more desirable way; for example, capro-lactam and/or sterols such as cholesterol, ergosterol, phytosterol, sitosterol, sitosterol pyroglutamate, 7-dehydro-cholesterol or lanosterol, may affect lipid films rigidity;

d) substances known to have antioxidant properties to improve the chemical stability of the components in the suspensions, such as tocopherol, propyl gallate, ascorbyl palmitate, or butylated hydroxy toluene.

The aqueous carrier in this invention is mostly water with possibly small quantities of physiologically compatible liquids such as isopropanol, glycerol, hexanol and the like (see for instance EP-A- 52.575). In general the amount of the organic hydrosoluble liquids will not exceed 5 - 10% by weight.

The present composition may also contain dissolved or suspended therein hydrophilic compounds and polymers defined generally under the name of viscosity enhancers or stabilizers. Although the presence of such compounds is not compulsory for ensuring stability to the air or gas bubbles with time in the present dispersions, they are advantageous to give some kind of "body" to the solutions. When desired, the upper concentrations of such additives when totally innocuous can be very high, for instance up to 80 - 90% by weight of solution with Iopamidol and other iodinated X-ray contrast agents. However with other viscosity enhancers like for instance sugars, e.g. lactose, sucrose, maltose, galactose, glucose, etc. or hydrophilic polymers like starch, dextran, polyvinyl alcohol, polyvinyl-pyrrolidone, dextrin, xanthan or partly hydrolyzed cellulose oligomers,

as well as proteins and polypeptides, the concentrations are best between about 1 and 40% by weight, a range of about 5 - 20% being preferred.

Like in the prior art, the injectable compositions of this invention can also contain physiologically acceptable electrolytes; an example is an isotonic solution of salt.

The present invention naturally also includes dry storable pulverulent blends which can generate the present microbubble containing dispersions upon simple admixing with water or an aqueous carrier phase. Preferably such dry blends or formulations will contain all solid ingredients necessary to provide the desired microbubbles suspensions upon the simple addition of water, i.e. principally the surfactants in lamellar form containing trapped or adsorbed therein the air or gas required for microbubble formation, and accessorially the other non-film forming surfactants, the viscosity enhancers and stabilizers and possibly other optional additives. As said before, the air or gas entrappment by the laminated surfactants occurs by simply exposing said surfactants to the air (or gas) at room or super-atmospheric pressure for a time sufficient to cause said air or gas to become entrapped within the surfactant. This period of time can be very short, e.g. in the order of a few seconds to a few minutes although over-exposure, i.e. storage under air or under a gaseous atmosphere is in no way harmful. What is important is that air can well contact as much as possible of the available surface of the laminated surfactant, i.e. the dry material should preferably be in a "fluffy" light flowing condition. This is precisely this condition which results from the freeze-drying of an aqueous solution of liposomes and hydrophilic agent as disclosed in US-A- 4,229,360.

In general, the weight ratio of surfactants to hydrophilic viscosity enhancer in the dry formulations will be in the order of 0.1:10 to 10:1, the further optional ingredients, if any, being present in a ratio not exceeding 50% relative to the total of surfactants plus viscosity enhancers.

The dry blend formulations of this invention can be prepared by very simple methods. As seen before, one preferred method is to first prepare an aqueous solution in which the film

forming lipids are laminarized, for instance by sonication, or using any conventional technique commonly used in the liposome field, this solution also containing the other desired additives, i.e. viscosity enhancers, non-film forming surfactants, electrolyte, etc., and thereafter freeze drying to a free flowable powder which is then stored in the presence of air or an entrappable gas.

The dry blend can be kept for any period of time in the dry state and sold as such. For putting it into use, i.e. for preparing a gas or air microbubble suspension for ultrasonic imaging, one simply dissolves a known weight of the dry pulverulent formulation in a sterile aqueous phase, e.g. water or a physiologically acceptable medium. The amount of powder will depend on the desired concentration of bubbles in the injectable product, a count of about  $10^8$  -  $10^9$  bubbles/ml being generally that from making a 5 - 20% by weight solution of the powder in water. But naturally this figure is only indicative, the amount of bubbles being essentially dependent on the amount of air or gas trapped during manufacture of the dry powder. The manufacturing steps being under control, the dissolution of the dry formulations will provide microbubble suspensions with well reproducible counts.

The resulting microbubble suspensions (bubble in the 0.5 - 10  $\mu$ m range) are extraordinarily stable with time, the count originally measured at start staying unchanged or only little changed for weeks and even months; the only observable change is a kind of segregation, the larger bubbles (around 10  $\mu$ m) tending to rise faster than the small ones.

It has also been found that the microbubbles suspensions of this invention can be diluted with very little loss in the number of microbubbles to be expected from dilution, i.e. even in the case of high dilution ratios, e.g.  $1/10^2$  to  $1/10^4$ , the microbubble count reduction accurately matches with the dilution ratio. This indicates that the stability of the bubbles depends on the surfactant in lamellar form rather than on the presence of stabilizers or viscosity enhancers like in the prior art. This property is advantageous in regard to imaging test reproducibility as the bubbles are not affected by dilution with

blood upon injection into a patient.

Another advantage of the bubbles of this invention versus the microcapsules of the prior art surrounded by a rigid but breakable membrane which may irreversibly fracture under stress is that when the present suspensions are subject to sudden pressure changes, the present bubbles will momentarily contract elastically and then resume their original shape when the pressure is released. This is important in clinical practice when the microbubbles are pumped through the heart and therefore are exposed to alternating pressure pulses.

The reasons why the microbubbles in this invention are so stable are not clearly understood. Since to prevent bubble escape the buoyancy forces should equilibrate with the retaining forces due to friction, i.e. to viscosity, it is theorized that the bubbles are probably surrounded by the laminated surfactant. Whether this laminar surfactant is in the form of a continuous or discontinuous membrane, or even as closed spheres attached to the microbubbles, is for the moment unknown but under investigation. However the lack of a detailed knowledge of the phenomena presently involved does not preclude full industrial operability of the present invention.

The bubble suspensions of the present invention are also useful in other medical/diagnostic applications where it is desirable to target the stabilized microbubbles to specific sites in the body following their injection, for instance to thrombi present in blood vessels, to atherosclerotic lesions (plaques) in arteries, to tumor cells, as well as for the diagnosis of altered surfaces of body cavities, e.g. ulceration sites in the stomach or tumors of the bladder. For this, one can bind monoclonal antibodies tailored by genetic engineering, antibody fragments or polypeptides designed to mimic antibodies, bioadhesive polymers, lectins and other site-recognizing molecules to the surfactant layer stabilizing the microbubbles. Thus monoclonal antibodies can be bound to phospholipid bilayers by the method described by L.D. Leserman, P. Machy and J. Barbet ("Liposome Technology vol. III" p. 29 ed. by G. Gregoriadis, CRC Press 1984). In another approach a palmitoyl antibody is first synthesized and then incorporated in phospholipid bilayers



following L. Huang, A. Huang and S.J. Kennel ("Liposome Technology vol. III" p. 51 ed. by G. Gregoriadis, CRC Press 1984). Alternatively, some of the phospholipids used in the present invention can be carefully selected in order to obtain preferential uptake in organs or tissues or increased half-life in blood. Thus GM1 gangliosides- or phosphatidylinositol-containing liposomes, preferably in addition to cholesterol, will lead to increased half-lives in blood after intravenous administration in analogy with A. Gabizon, D. Papahadjopoulos, Proc. Natl Acad. Sci USA 85 (1988) 6949.

The gases in the microbubbles of the present invention can include, in addition to current innocuous physiologically acceptable gases like CO<sub>2</sub>, nitrogen, N<sub>2</sub>O, methane, butane, freon and mixtures thereof, radioactive gases such as <sup>133</sup>Xe or <sup>81</sup>Kr are of particular interest in nuclear medicine for blood circulation measurements, for lung scintigraphy etc.

The following Examples illustrate the invention on a practical stand point.

#### Echogenic measurements

Echogenicity measurements were performed in a pulse - echo system made of a plexiglas specimen holder (diameter 30 mm) and a transducer holder immersed in a constant temperature water bath, a pulser-receiver (Accutron M3010S) with for the receiving part an external pre-amplifier with a fixed gain of 40 dB and an internal amplifier with adjustable gain from -40 to +40 dB. A 10MHz low-pass filter was inserted in the receiving part to improve the signal to noise ratio. The A/D board in the IBM PC was a Sonotek STR 832. Measurements were carried out at 2.25, 3.5, 5 and 7.5 MHz.

#### Example 1

A liposome solution (50 mg lipids per ml) was prepared in distilled water by the REV method (see F. Szoka Jr. and D. Papahadjopoulos, Proc. Natl. Acad. Sci. USA 75 (1978) 4194) using hydrogenated soya lecithin (NC 95 H, Nattermann Chemie, Köln, W. Germany) and dicetylphosphate in a molar ratio 9/1. This liposome preparation was extruded at 65°C (to calibrate the

vesicle size) through a 1  $\mu\text{m}$  polycarbonate filter (Nucleopore). Two ml of this solution were admixed with 5 ml of a 75% iopamidol solution in water and 0.4 ml of air and the mixture was forced back and forth through a two syringe system as disclosed in DE-A-3529195, while maintaining continuously a slight over-pressure. This resulted in the formation of a suspension of microbubbles of air in the liquid ( $10^5$ - $10^6$  bubbles per ml, bubble size 1-20  $\mu\text{m}$  as estimated by light microscopy) which was stable for several hours at room temperature. This suspension gave a strong echo signal when tested by ultrasonic echography at 7.5, 5, 3.5 and 2.25 MHz.

#### Example 2

A distilled water solution (100 ml) containing by weight 2% of hydrogenated soya lecithin and dicetylphosphate in a 9/1 molar ratio was sonicated for 15 min at 60-65°C with a Branson probe sonifier (Type 250).

After cooling, the solution was centrifuged for 15 min at 10,000 g and the supernatant was recovered and lactose added to make a 7.5% b.w. solution. The solution was placed in a tight container in which a pressure of 4 bar of nitrogen was established for a few minutes while shaking the container. Afterwards, the pressure was released suddenly whereby a highly concentrated bubble suspension was obtained ( $10^{10}$  -  $10^{11}$  bubbles/ml). The size distribution of the bubbles was however wider than in Example 1, i.e. from about 1 to 50  $\mu\text{m}$ . The suspension was very stable but after a few days a segregation occurred in the standing phase, the larger bubbles tending to concentrate in the upper layers of the suspension.

#### Example 3

Twenty g of glass beads (diameter about 1 mm) were immersed into a solution of 100 mg of dipalmitoylphosphatidylcholine (Fluka A.G. Buchs) in 10 ml of chloroform. The beads were rotated under reduced pressure in a rotating evaporator until all  $\text{CHCl}_3$  had escaped. Then the beads were further rotated under atmospheric pressure for a few minutes and 10 ml of distilled water were added. The beads were removed and a suspension of air

microbubbles was obtained which was shown to contain about  $10^6$  bubbles/ml after examination under the microscope. The average size of the bubbles was about 3 - 5  $\mu\text{m}$ . The suspension was stable for several days at least.

#### Example 4

A hydrogenated soya lecithin/dicetylphosphate suspension in water was laminarized using the REV technique as described in Example 1. Two ml of the liposome preparation were added to 8 ml of 15% maltose solution in distilled water. The resulting solution was frozen at  $-30^\circ\text{C}$ , then lyophilized under 0.1 Torr. Complete sublimation of the ice was obtained in a few hours. Thereafter, air pressure was restored in the evacuated container so that the lyophilized powder became saturated with air in a few minutes.

The dry powder was then dissolved in 10 ml of sterile water under gentle mixing, whereby a microbubble suspension ( $10^8$ - $10^9$  microbubbles per ml, dynamic viscosity  $< 20 \text{ mPa.s}$ ) was obtained. This suspension containing mostly bubbles in the 1-5  $\mu\text{m}$  range was stable for a very long period, as numerous bubbles could still be detected after 2 months standing. This microbubble suspension gave a strong response in ultrasonic echography. If in this example the solution is frozen by spraying in air at  $-30$  to  $-70^\circ\text{C}$  to obtain a frozen snow instead of a monolithic block and the snow is then evaporated under vacuum, excellent results are obtained.

#### Example 5

Two ml samples of the liposome solution obtained as described in Example 4 were mixed with 10 ml of an 5% aqueous solution of gelatin (sample 5A), human albumin (sample 5B), dextran (sample 5C) and iopamidol (sample 5D). All samples were lyophilized. After lyophilization and introduction of air, the various samples were gently mixed with 20 ml of sterile water. In all cases, the bubble concentration was above  $10^8$  bubbles per ml and almost all bubbles were below 10  $\mu\text{m}$ . The procedure of the foregoing Example was repeated with 9 ml of the liposome preparation (450 mg of lipids) and only one ml of a 5% human

albumin solution. After lyophilization, exposure to air and addition of sterile water (20 ml), the resulting solution contained  $2 \times 10^8$  bubbles per ml, most of them below 10  $\mu\text{m}$ .

#### Example 6

Lactose (500 mg), finely milled to a particle size of 1-3  $\mu\text{m}$ , was moistened with a chloroform (5 ml) solution of 100 mg of dimyristoylphosphatidylcholine/cholesterol/dipalmitoylphosphatidic acid (from Fluka) in a molar ratio of 4:1:1 and thereafter evaporated under vacuum in a rotating evaporator. The resulting free flowing white powder was rotated a few minutes under nitrogen at normal pressure and thereafter dissolved in 20 ml of sterile water. A microbubble suspension was obtained with about  $10^5$ - $10^6$  microbubbles per ml in the 1-10  $\mu\text{m}$  size range as ascertained by observation under the microscope. In this Example, the weight ratio of coated surfactant to water-soluble carrier was 1:5. Excellent results ( $10^7$ - $10^8$  microbubbles/ml) are also obtained when reducing this ratio to lower values, i.e. down to 1:20, which will actually increase the surfactant efficiency for the intake of air, that is, this will decrease the weight of surfactant necessary for producing the same bubble count.

#### Example 7

An aqueous solution containing 2% of hydrogenated soya lecithin and 0.4% of Pluronic<sup>®</sup> F68 (a non ionic polyoxyethylene-polyoxypropylene copolymer surfactant) was sonicated as described in Example 2. After cooling and centrifugation, 5 ml of this solution were added to 5 ml of a 15% maltose solution in water. The resulting solution was frozen at  $-30^\circ\text{C}$  and evaporated under 0.1 Torr. Then air pressure was restored in the vessel containing the dry powder. This was left to stand in air for a few seconds, after which it was used to make a 10% by weight aqueous solution which showed under the microscope to be a suspension of very tiny bubbles (below 10  $\mu\text{m}$ ); the bubble concentration was in the range of  $10^7$  bubbles per ml. This preparation gave a very strong response in ultrasonic echography at 2.25, 3.5, 5 and 7.5 MHz.

Example 8

Two-dimensional echocardiography was performed in an experimental dog following peripheral vein injection of 0.1-2 ml of the preparation obtained in Example 4. Opacification of the left heart with clear outlining of the endocardium was observed, thereby confirming that the microbubbles (or at least a significant part of them) were able to cross the pulmonary capillary circulation.

Example 9

A phospholipid/maltose lyophilized powder was prepared as described in Example 4. However, at the end of the lyophilization step, a  $^{133}\text{Xe}$  containing gas mixture was introduced in the evacuated container instead of air. A few minutes later, sterile water was introduced and after gentle mixing a microbubble suspension containing  $^{133}\text{Xe}$  in the gas phase was produced. This microbubble suspension was injected into living bodies to undertake investigations requiring use of  $^{133}\text{Xe}$  as tracer. Excellent results were obtained.

CLAIMS

1. A composition adapted for injection into the bloodstream and body cavities of living beings, e.g. for the purpose of ultrasonic echography, consisting of a suspension of air or gas microbubbles in a physiologically acceptable aqueous carrier phase comprising from about 0.01 to about 20% by weight of one or more dissolved or dispersed surfactants, characterized in that at least one of the surfactants is a film forming surfactant present in the composition at least partially in lamellar or laminar form.

2. The composition of claim 1, characterized in that the lamellar surfactant is in the form of mono- or pluri-molecular membrane layers.

3. The composition of claim 1, characterized in that the lamellar surfactant is in the form of liposome vesicles.

4. The composition of claim 1, characterized in that it essentially consists of a liposome solution containing air or gas microbubbles developed therein.

5. The composition of claim 4, characterized in that the size of most of both liposomes and microbubbles is below 50  $\mu\text{m}$ , preferably below 10  $\mu\text{m}$ .

6. The composition of claim 1, containing about  $10^8 - 10^9$  bubbles of 0.5 - 10  $\mu\text{m}$  size/ml, said concentration showing little or substantially no variability under storage for at least a month.

7. The composition of claim 1, characterized in that the surfactants are selected from phospholipids including the lecithins such as phosphatidic acid, phosphatidyl-choline, phosphatidyl-ethanolamine, phosphatidyl-serine, phosphatidyl-glycerol phosphatidyl-inositol, cardiolipin and sphingomyelin.

8. The composition of claim 7, characterized in further containing substances affecting the properties of liposomes selected from phosphatidyl-glycerol, dicetyl-phosphate, cholesterol, ergosterol, phytosterol, sitosterol, lanosterol, tocopherol, propyl gallate, ascorbyl palmitate and butylated hydroxytoluene.

9. The composition of claim 1, further containing dissolved

viscosity enhancers or stabilizers selected from linear and cross-linked poly- and oligo-saccharides, sugars, hydrophilic polymers and iodinated compounds such as Iopamidol in a weight ratio to the surfactants comprised between about 1:5 to 100:1.

10. The composition of claim 1, in which the surfactants comprise up to 50% by weight of non-laminar surfactants selected from fatty acids, esters and ethers of fatty acids and alcohols with polyols such as polyalkylene glycols, polyalkylenated sugars and other carbohydrates, and polyalkylenated glycerol.

11. A method for the preparation of the suspensions of claim 1, characterized by the following steps:

(a) selecting at least one film forming surfactant and converting it into lamellar form;

(b) contacting the surfactant in lamellar form with air or an adsorbable or entrappable gas for a time sufficient for that air or gas to become bound by said surfactant; and

(c) admixing the surfactant in lamellar form with an aqueous liquid carrier, whereby a stable dispersion of air or gas microbubbles in said liquid carrier will result.

12. The method of claim 11, in which step (c) is brought about before step (b), the latter being effected by introducing pressurized air or gas into the liquid carrier and thereafter releasing the pressure.

13. The method of claim 11, in which step (c) is brought about by gentle mixing of the components, no shaking being necessary, whereby the air or gas bound to the lamellar surfactant in step (b) will develop into a suspension of stable microbubbles.

14. The method of claims 11 or 12, in which the liquid carrier contains dissolved therein stabilizer compounds selected from hydrosoluble proteins, polypeptides, sugars, poly- and oligo-saccharides and hydrophilic polymers.

15. The method of claim 11, in which the conversion of step (a) is effected by coating the surfactant onto particles of soluble or insoluble materials; step (b) is effected by letting the coated particles stand for a while under air or a gas; and step (c) is effected by admixing the coated particles with an aqueous liquid carrier.

16. The method of claim 11, in which the conversion of step (a) is effected by sonicating or homogenizing under high pressure an aqueous solution of film forming lipids, this operation leading, at least partly, to the formation of liposomes.

17. The method of claim 16, in which step (b) is effected by freeze-drying the liposome containing solution, the latter optionally containing hydrophilic stabilizers and contacting the resulting freeze-dried product with air or a gas for a period of time.

18. The method of claims 16 and 17, in which the water solution of film forming lipids also contains viscosity enhancers or stabilizers selected from hydrophilic polymers and carbohydrates in weight ratio relative to the lipids comprised between 1:5 and 100:1.

19. A dry pulverulent formulation which, upon dissolution in water, will form an aqueous suspension of microbubbles for ultrasonic echography, characterized in containing one or more film forming surfactants in laminar form and hydrosoluble stabilizers.

20. The dry formulation of claim 19, in which the surfactants in laminar form are in the form of fine layers deposited on the surface of soluble or insoluble solid particulate material.

21. The dry formulation of claim 20, in which the insoluble solid particles are glass or polymer beads.

22. The dry formulation of claim 20, in which the soluble particles are made of hydrosoluble carbohydrates, polysaccharides, synthetic polymers, albumin, gelatin or Iopamidol.

23. The dry formulation of claim 19, which comprises freeze-dried liposomes.

24. The use of the injectable composition of claim 1 for ultrasonic echography.

25. The use of the injectable composition of claims 1-10 for transporting in the blood-stream or body cavities bubbles of foreign gases active therapeutically or diagnostically.

26. The composition of claim 4, in which the surfactant comprises, bound thereto, bioactive species designed for



specific targeting purposes, e.g. for immobilizing the bubbles in specifically defined sites in the circulatory system, or in organs, or in tissues.

27. The composition of claim 4, in which the surfactant comprises, bound thereto, bioactive species selected from monoclonal antibodies, antibody fragments or polypeptides designed to mimic antibodies, bioadhesive polymers, lectins and other receptor recognizing molecules.

ABSTRACT

Gas or air filled microbubble suspensions in aqueous phases usable as imaging contrast agents in ultrasonic echography. They contain laminarized surfactants and, optionally, hydrophilic stabilizers. The laminarized surfactants can be in the form of liposomes. The suspensions are obtained by exposing the laminarized surfactants to air or a gas before or after admixing with an aqueous phase.

01/775989 #3

LARRY S. NIXON NIXON & VANDERHYE P.C. 2200 CLARENDON BOULEVARD, 14TH FLOOR ARLINGTON, VIRGINIA 22201		UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) NOTIFICATION OF ACCEPTANCE OF APPLICATION UNDER 35 U.S.C. 371 AND 37 CFR 1.494 OR 1.495	
		Date of Mailing <b>02 MAR 1992</b>	
		File Reference 1201-3	
IDENTIFICATION OF THE INTERNATIONAL APPLICATION			
International application Number PCT/EP91/00620	International filing date 02 APRIL 1991	Priority date claimed 02 APRIL 1990	
Applicant for DO/EO/US SCHNEIDER ET AL.			
NOTIFICATION			
The applicant is hereby advised that the United States Patent and Trademark Office in its capacity as a <input checked="" type="checkbox"/> Designated Office, <input type="checkbox"/> Elected Office, has determined that the above identified international application has met the requirements of 35 U.S.C. 371 and 37 CFR <input checked="" type="checkbox"/> 1.494, <input type="checkbox"/> 1.495 and is ACCEPTED for national patentability examination in the United States Patent and Trademark Office.			
The United States Serial Number assigned to the application and the relevant dates are: <u>01/775989</u> <u>20 NOV 1991</u> <u>20 NOV 1991</u>			
U.S. NATIONAL SERIAL NO.    35 U.S.C. 102(e) DATE    DATE OF RECEIPT 35 U.S.C. 371 REQUIREMENTS			
<input type="checkbox"/> A request for immediate examination under 35 U.S.C. 371 (f) was received on <u>20 NOV 1991</u> and the application will be examined in turn.			
<input type="checkbox"/> No request for immediate examination under 35 U.S.C. 371(f) was received. The application will not be processed or examined before the time limit set forth in either			
<input type="checkbox"/> PCT Article 23 (Chapter I of the PCT), or <input type="checkbox"/> PCT Article 40 (Chapter II of the PCT) whichever is applicable.			
UNITED STATES DESIGNATED/ ELECTED OFFICE			
ADDRESS ONLY: COMMISSIONER OF PATENTS AND TRADEMARKS Box PCT, Attn. DO/EO/US Washington, D.C. 20231		AUTHORIZED OFFICER PAUL J. KIDWELL <i>P. Kidwell</i> INTL. COMM. ON PAT. CO-OPERATION	

9 DEC 1991

PATENT COOPERATION TREATY

COMMUNICATION OF  
INTERNATIONAL APPLICATIONS  
under PCT Article 20

DATE OF MAILING  
14 November 1991 (14.11.91)

To: United States Patent  
and Trademark Office  
Washington, D.C.  
  
in its capacity as a designated Office

From : The International Bureau of WIPO  
1211 Geneva 20  
Switzerland

Pursuant to PCT Article 20 and PCT Rule 47, the International Bureau herewith transmits copies of the international applications having the following international application and international publication numbers:

INT. APPL. NO.

INT. PUBL. NO.

INT. APPL. NO.

INT. PUB. NO.

PCT/EP91/00620

W091/15244

CORRECTED VERSION